

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 9/46, A23L 2/40, 1/236, A61K 31/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/20077</b> <b>(43) International Publication Date:</b> 15 September 1994 (15.09.94)
<b>(21) International Application Number:</b> PCT/HU94/00006 <b>(22) International Filing Date:</b> 10 March 1994 (10.03.94) <b>(30) Priority Data:</b> P 93 00680 10 March 1993 (10.03.93) HU <b>(71) Applicant (for all designated States except US):</b> BÉRES RÉSZVÉNYTÁRSASÁG [HU/HU]; Szépvölgyi út 135, H-1037 Budapest (HU). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BÉRES, József [HU/HU]; Baross G. u. 3, H-4600 kisvárd (HU). BÉRES, József [HU/HU]; Törökvész u. 74/b, H-1025 Budapest (HU). LEX, László [HU/HU]; Kéve u. 4, H-1033 Budapest (HU). BÁRKÁNYI, Istvánné [HU/HU]; Élmunkás u. 1, H-4440 Tiszavasvári (HU). <b>(74) Agent:</b> DANUBIA PATENT & TRADEMARK ATTORNEYS; Bajcsy-Zsilinszky út 16., H-1051 Budapest (HU).	<b>(81) Designated States:</b> AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> SUGAR- AND SODIUM-FREE EFFERVESCENT TABLETS AND GRANULES AND PROCESS FOR PREPARING SAME  <b>(57) Abstract</b>  The invention relates to effervescent tablets and granules comprising a shell material, a basic sparkling component, an acidic sparkling component, and a sweetening agent, furthermore macro- and microelements and optionally vitamins as active agents. The effervescent tablets and granules comprise 20-50 % by mass of mannitol as shell material, 8-25 % by mass of potassium hydrogen carbonate as basic sparkling component, 9-27 % by mass of malic acid as acidic sparkling component, and 0.4-2.2 % by mass of aspartame as sweetening agent. Furthermore, the invention relates to a process for preparing the above-described effervescent tablets and granules.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MD	Republic of Moldova	TJ	Tajikistan
DE	Germany	MG	Madagascar	TT	Trinidad and Tobago
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain	MN	Mongolia	US	United States of America
FI	Finland			UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**Sugar- and sodium-free effervescent tablets and granules  
and process for preparing same**

The invention relates to sugar- and sodium-free  
5 effervescent tablets and granules and a process for their  
preparation.

More specifically, the invention relates to effervescent  
tablets and granules comprising a shell material, a basic  
effervescent and disintegrating (further on: sparkling) com-  
10 ponent, an acidic sparkling component, and a sweetening  
agent, further macro- and trace elements and optionally vit-  
amins. Furthermore, the invention relates to a process for  
the preparation of such tablets and granules.

It is known that nowadays one of the most popular phar-  
15 maceutical form for introducing medicines, vitamins and  
mineral substances to the organism is the so-called efferv-  
escent tablet (Pharmaceutical Dosage Form: Tablets, Vol.  
I., 2nd edition, A. Lieberman ed., 1989, Marcel Dekker,  
Inc.). The development of this form is supported from the  
20 viewpoint of pharmaceutical effect, beside commercial  
reasons, by several factors, e.g. the decrease of stomach  
irritation, the enhancement of absorption, etc. The solution  
of such tablets in water results in a carbonated or sparkl-  
ing beverage containing carbon dioxide.

25 The spectacular disintegration of effervescent tablets  
is caused by a mixture consisting of an acid and a base,  
wherein said mixture when contacting with water nearly  
wedges the tablet during evolution of carbon dioxide.

The manufacturing and packaging of effervescent tablets  
30 requires high care; consequently, in the practice the direct  
compressing method is preferred to the wet processes.

Most effervescent tablets comprise three main components  
beside the active agents: a shell and binding agent, an  
acidic sparkling agent and a basic sparkling agent.

35 As shell and binding agent sugars (lactose, saccharose,

- 2 -

glucose), sorbitol, xylitol or starch, as acidic sparkling agent citric acid, tartaric acid, fumaric acid or adipic acid while as basic sparkling component sodium hydrogen carbonate, sodium carbonate and magnesium carbonate are generally used.

From the other components generally used in effervescent tablets the following preferable agents are mentioned: sweetening agents such as sugars, saccharin, sodium cyclamate and aspartame; flavouring and aroma agents; lubricating agents such as polyethylene glycols, silicone oils, stearates and adipic acid.

The U.S. patent specification No. 4,725,427 describes an effervescent tablet which contains lactose as shell material, citric acid as acidic sparkling agent, a mixture of sodium hydrogen carbonate and potassium hydrogen carbonate as basic sparkling component and aspartame as sweetening agent. Beside water and fat-soluble vitamins this tablet contains inorganic substances as active agents and these substances are rendered more utilizable biologically by using them in chelate form. This composition does not render possible, however, the preparation of sodium-free tablets and this is a considerable disadvantage since it is well-known that the introduction of an excess of sodium into the organism results in several deleterious physiological effects. A further disadvantage of this composition is caused by the presence of citric acid in an amount of 20-45 % by mass since the high amount of this acid may result in harmful physiological effects.

The U.S. patent specification No. 4,678,661 describes effervescent tablets containing a mixture of calcium carbonate and potassium carbonate as basic sparkling component. It is a considerable disadvantage of this tablet that the disagreeable soapy flavour of potassium hydrogen carbonate can be tasted in it. Besides, the use of calcium carbonate unfavourably influences the dissolution time.

The U.S. patent specification No. 4,704,269 describes an effervescent tablet comprising potassium hydrogen carbonate

- 3 -

as basic sparkling component, malic acid and citric acid as acidic sparkling components, a mixture of sorbitol and maltodextrin as shell and binding agents and calcium saccharate as sweetening agent. This composition is used as antacid and  
5 analgesic; its disadvantage is that, owing to the presence of sorbitol, its storability is not satisfactory. Besides, the sorbitol is not recommended for general use in soft drinks since some individuals have a low gastric tolerance for it (Martindale: The Extra Pharmacopoeia, 19th ed.,  
10 London, 1989, p. 1274).

The invention aims at preparing effervescent tablets and granules which are chemically stable, can readily be compressed, have advantageous physical properties, are free from sodium and sugar, and contain macro- and trace elements  
15 and optionally vitamins in a homogeneous distribution.

The invention is based on the recognition that the above aim can be completely attained if the following components are used as basic materials for preparing effervescent tablets and granules: mannitol as shell material, malic acid  
20 as acidic sparkling component, potassium hydrogen carbonate as basic sparkling component and aspartame as sweetening agent.

The invention is further based on the recognition that the use of mannitol renders possible the introduction of  
25 salts of macro- and trace elements of high crystal-water content into the composition. Consequently, the invention is based on the overcoming of a technical prejudice since until now it was known that no effervescent tablets and granules can be prepared from such materials due to the fact that the  
30 high water content inhibits the compressing and at the same time it results in the premature deliquesce of the tablets.

A further basis of the invention is the recognition that in case of using mannitol for preparing tablets or granules the macro- and trace elements form complexes with the mannitol whereby the incompatibilities of the components can be  
35 eliminated during the technological processes, the end-product will be chemically stable, and the mannitol complexes

- 4 -

obtained can be more readily absorbed by the organism, that is, they become better utilizable.

A further basis of the invention is the recognition that, in the case of the common use of mannitol, malic acid  
5 and aspartame, the potassium hydrogen carbonate can be used even alone as basic sparkling component whereby it is possible to eliminate the sodium ions from the composition. Besides, with such a combination the unagreeable flavour and the sensitivity to moisture of potassium hydrogen carbonate  
10 can be eliminated. Furthermore, in the case of this composition the bad compressibility of potassium hydrogen carbonate, i.e. the property that it strongly adheres to the surface of stamps and matrices which renders impossible its compressing in spaces containing a relative moisture content  
15 of 45 % or more, can be eliminated. Consequently, the invention is based even in this respect on the overcoming of a technical prejudice. This statement is proved by the fact that the US patent specification No. 4,678,661 contains in column 1, lines 27-32 the following statement: "The utilization of potassium bicarbonate and potassium carbonate alone  
20 fails to meet the need because, first, the potassium compounds give the substance an unpleasant soapy taste, and second, the moisture sensitivity due to the introduction of potassium salts leads to great technical problems."

25 A further basis of the invention is the recognition that in case of the common use of malic acid as acidic sparkling component with mannitol a suitably compressible composition is obtained. This recognition is surprising since it is known that the malic acid alone can not be readily compressed and due to its low melting point it is a technologically  
30 difficultly treatable compound which melts while grinding. On the other hand, the use of malic acid in a relatively high amount is rendered possible by our recognition and at the same time the antioxidant and flavour-improving effect of malic acid as well as its ability to opti-  
35 mally setting the pH value can be utilized.

Finally, the invention is based on the recognition that

- 5 -

by the common use of mannitol, potassium hydrogen carbonate, malic acid and aspartame it is rendered possible to prepare a composition of low energy content which does not cause gastrointestinal complaints. The tablets prepared from this  
5 composition have very high breaking strength and result in a quickly sparkling and clear solution although the composition comprises incompatible vitamins, macro- and trace elements and components (potassium hydrogen carbonate, malic acid, salts of macro- and trace elements of high crystal-  
10 water content) having *per se* bad compressing properties.

Based on the above the invention relates to effervescent tablets and granules comprising a shell material, a basic sparkling component, an acidic sparkling component, and a sweetening agent, furthermore macro- and microelements and  
15 optionally vitamins as active agents. According to the invention the effervescent tablets and granules comprise 20-50 % by mass, preferably 30-40 % by mass, of mannitol as shell material, 8-25 % by mass, preferably 14-18 % by mass, of potassium hydrogen carbonate as basic sparkling compo-  
20 nent, 9-27 % by mass, preferably 15-21 % by mass, of malic acid as acidic sparkling component, and 0.4-2.2 % by mass, preferably 0.6-1.5 % by mass, of aspartame as sweetening agent, furthermore, if desired, flavouring, lubricating and other additives generally used in the manufacture of efferver-  
25 vescent tablets in an amount necessary to supplement the mass of the components to 100 %.

Furthermore, the invention relates to a process for preparing effervescent tablets or granules. According to the invention one proceeds by homogenizing 20-50 % by mass, preferably 30-40 % by mass, of mannitol, 8-25 % by mass, preferably 14-18 % by mass, of potassium hydrogen carbonate, 9-24 % by mass, preferably 15-21 % by mass, of malic acid, and 0.4-2.2 % by mass, preferably 0.6-1.5 % by mass, of aspartame together with the macro- and trace elements and vit-  
30 amins to be introduced and optionally together with flavouring, lubricating and other additives generally used in the manufacture of effervescent tablets, then granulating the

- 6 -

thus-obtained homogenizate to granules ready for compressing, and finally compressing tablets or granules of the desired size and strength.

The effervescent tablets and granules according to the invention contain as macro- and trace elements preferably magnesium, zinc, iron(II), copper(II), manganese(II), and chromium(III) cations, further molybdenum(VI) and selenium(IV) anions.

The iron ions are preferably used in the composition in the form of iron(II)-sulphate heptahydrate, the zinc ions in the form of zinc sulphate heptahydrate, the copper ions in the form of copper sulphate pentahydrate, the manganese ions in the form of manganese sulphate monohydrate, the molybdenum ions in the form of ammonium heptamolybdenate tetrahydrate, the selenium ions in the form of selenious acid, the magnesium ions in the form of magnesium sulphate heptahydrate, and the chromium ions in the form of chromium(III) chloride hexahydrate.

The vitamins are added to the composition preferably in the following amounts: 0.01-0.5 % by mass of vitamin B<sub>1</sub>, 0.01-0.25 % by mass of vitamin B<sub>2</sub>, 0.01-0.5 % by mass of vitamin B<sub>6</sub>, 0.001-0.01 % by mass of vitamin B<sub>12</sub>, 0.1-2 % by mass of nicotinamide, 0.01-0.5 % by mass of vitamin A, 0.0015-0.015 % by mass of vitamin D, 0.1-5 % by mass of vitamin C, 0.01-0.1 % by mass of folic acid, 0.1-0.5 % by mass of pantothenic acid, 0.01-7 % by mass of vitamin E and 0.001-0.01 % by mass of vitamin H.

The tablets according to the invention may contain besides the macro- and trace elements and vitamins also flavouring and aroma agents such as orange, lemon or pineapple aroma, lubricating agents such as polyethylene glycols, silicone oils, stearates or adipic acid, agents enhancing absorption such as tartaric acid and glycine, further any other additive usual in the manufacture of effervescent tablets.

The main advantages of the invention are as follows:

a) The tablets are chemically stable, can be readily com-



- 7 -

pressed and have excellent physical properties.

b) The tablets and granules contain the active agents, that is the macro- and trace elements as well as the vitamins, in homogeneous distribution.

5 c) The tablets give after dissolving in water a clear beverage of agreeable flavour, without any sediment.

d) The presence of mannitol renders possible the use of malic acid as acidic sparkling component in a relatively high amount whereby the agreeable antioxidant, flavour-im-  
10 proving and optimal pH-setting effects of this acid can be enforced.

e) By using the mannitol effervescent tablets of low calory content and rich in macro- and trace elements and vitamins can be prepared which can be consumed also by diabetic  
15 persons.

f) In the known effervescent tablets containing vitamins and mineral agents the trace elements are used in crystal-water-free form or in a form containing only a small amount of crystal water. On the other hand, the invention renders  
20 possible the use of substances of high crystal-water content, which *per se* are only badly compressible or cannot be compressed at all, are the most stable modifications of the inorganic compounds and thereby can be prepared or procured at a lower price and in high purity.

25 g) By the common use of mannitol, malic acid and aspartame the homogeneous distribution of the macro- and trace elements and vitamins can be realized even when their amount is very low in relation to the mass of the ready tablet. The homogeneous distribution of vitamins is ensured without disadvantageously influencing the properties of these sensitive  
30 compounds during the technological procedures.

h) The invention renders possible to prepare effervescent tablets containing incompatible active agents such as vitamins as well as macro- and trace elements.

35 i) During the manufacture of the tablets the macro- and trace elements form complexes with the mannitol, whereby the chemical stability of the tablet as well as the absorption

- 8 -

and biological utilization of the active agents become more favourable.

j) The invention renders possible the preparation of tablets by using sparkling components (potassium hydrogen carbonate and malic acid) and inorganic compounds with high crystal-water content (macro- and trace element sources) which, owing to their unfavourable properties, could not be previously used in the manufacture of effervescent tablets. Besides, the thus-obtained effervescent tablets have high breaking strength and give a clear solution with a short sparkling time.

The invention is further elucidated by the aid of the following non-limiting examples.

#### Example 1

The granules ready for pressing are composed of four granules and a so-called outer phase.

#### Granule I

	Vitamin B <sub>1</sub>	7.29	g
	Vitamin B <sub>2</sub>	7.50	g
20	Vitamin B <sub>6</sub>	10.94	g
	Ca-pantothenate	38.215	g
	Nicotinamide	85.00	g
	Mannitol	500.00	g

After sieving the materials are homogenized, kneaded with ethanol, granulated, then the wet granules are dried and re-granulated.

#### Granule II

30	Iron(II)-sulphate heptahydrate	99.55	g
	Malic acid	1500.00	g
	Mannitol	1500.00	g

After sieving the materials are homogenized, kneaded with distilled water, granulated, dried, then re-granulated and post-dried.

#### Granule III

35	Potassium hydrogen carbonate	3800.00	g
	Mannitol	3800.00	g

After sieving and homogenizing the mass is kneaded with a

- 9 -

mixture of water and ethanol, then after drying it is re-granulated.

5	<b>Granule IV</b>		
	Mannitol	3925.00	g
	Magnesium sulphate heptahydrate	1571.50	g
	Glycine	150.00	g
	Succinic acid	250.00	g
10	Mannitol	75.00	g
	Selenious acid	0.1635	g
	Ammonium heptamolybdenate tetrahydrate	0.690	g
	Manganese(II)-sulphate monohydrate	15.38	g
	Copper(II)-sulphate pentahydrate	29.47	g
15	Zinc sulphate heptahydrate	219.95	g

After grinding, homogenizing and kneading the mass is granulated with distilled water, then it is dried, re-granulated and post-dried.

20	<b>Materials of the outer phase</b>		
	Vitamine C	300.00	g
	Malic acid	3000.00	g
	Polyethylene glycol	710.00	g
	Aspartame	200.00	g
25	Lemon aroma	1000.00	g

After sieving and grinding the materials of the outer phase are homogenized. The thus-obtained pre-homogenizate is admixed with the granules I, II, III and IV and the thus-obtained mixture is homogenized. From the granules obtained in this way about 5000 tablets with a diameter of 32 mm and an average weight of 4.5 g are compressed.

### 35 Example 2

One proceeds in the same way as described in Example 1 with the difference that the vitamins are supplemented with vitamin E and the amounts of the components are changed in the following way:

- 10 -

	<u>Name of the component</u>	<u>Amount (g)</u>
	Iron(II)-sulphate ( $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ )	99.56
	Zinc(II)-sulphate ( $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ )	109.97
	Copper(II)-sulphate ( $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ )	14.74
5	Manganese(II)-sulphate ( $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ )	7.69
	Ammonium molybdenate [ $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ ]	0.276
	Selenious acid ( $\text{H}_2\text{SeO}_3$ )	0.082
	Magnesium sulphate ( $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ )	608.34
10	Vitamin B <sub>1</sub> (thiamine·HCl)	3
	Vitamin B <sub>2</sub> (riboflavine)	3.5
	Vitamin B <sub>6</sub> (pyridoxine·HCl)	4
	Nicotinamide	40
	Vitamin C [L-(+)-ascorbic acid]	175
15	Phantothenic acid (Ca-pantothenate)	15
	Vitamin E (DL- $\alpha$ -tocoferol)	25
	Succinic acid	100
	Glycine	75
20	Malic acid	2750
	Potassium hydrogen carbonate ( $\text{KHCO}_3$ )	2300
	Mannitol	6500
	Aspartame	200
	Pineapple aroma	1000
25	Polyethylene glycol	750

From the granules ready for pressing about 5000 tablets having a diameter of 25 mm and an average weight of 3 g are stamped.

### 30 Example 3

One proceeds as described in Example 1 with the difference that the microelements are supplemented with chromium and the vitamins with vitamins B<sub>12</sub>, A, D and H and folic acid, furthermore the amounts of the components are changed

35 in the following way:

<u>Name of the component</u>	<u>Amount (g)</u>
Iron(II)-sulphate ( $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ )	373.35

- 11 -

	Zinc(II)-sulphate ( $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ )	329.97
	Copper(II)-sulphate ( $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ )	39.29
	Manganese(II)-sulphate ( $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ )	38.46
	Ammonium molybdenate [ $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ ]	1.38
5	Selenious acid ( $\text{H}_2\text{SeO}_3$ )	0.2
	Magnesium sulphate ( $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ )	5069.5
	Chromium(III)-chloride ( $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ )	1.28
	Vitamin B <sub>1</sub> (thiamine·HCl)	7.5
10	Vitamin B <sub>2</sub> (riboflavine)	8.5
	Vitamin B <sub>6</sub> (pyridoxine·HCl)	10
	Vitamin B <sub>12</sub> (cyanocobalamin)	0.01
	Nicotinamide	95
	Vitamin A	5
15	Vitamin D	0.05
	Vitamin C [L-(+)-ascorbic acid]	450
	Folic acid	1
	Pantothenic acid (Ca-pantothenate)	35
	Vitamin E (DL- $\alpha$ -tocoferol)	50
20	Vitamin H (biotine)	325
	Succinic acid	300
	Glycine	180
	Malic acid	6000
25	Potassium hydrogen carbonate ( $\text{KHCO}_3$ )	5000
	Mannitol	11500
	Aspartame	300
	Orange aroma	1500
	Polyethylene glycol	2000
30	From the granules ready for compressing about 5000 tablets having a diameter of 35 mm and a weight of 6.6 g are compressed.	

#### Example 4

One proceeds as described in Example 3 with the difference that the amount of malic acid is changed to 3500 g, that of potassium hydrogen carbonate to 2800 g, that of mannitol to 16,000 g and that of aspartame to 150 g. From

- 12 -

the granules ready for compressing about 5000 tablets having a diameter of 32 mm and an average weight of 6.6 g are compressed.

#### Example 5

- 5        One proceeds as described in Example 3 with the difference that the amount of malic acid is changed to 10,000 g, that of potassium hydrogen carbonate to 9000 g, that of mannitol to 8000 g and that of aspartame to 800 g. From the granules ready for compressing about 5000 tablets having a
- 10   diameter of 32 mm and an average weight of 7.7 g are compressed.

What we claim is:

1. Effervescent tablets and granules comprising a shell material, a basic sparkling component, an acidic sparkling component and a sweetening agent, furthermore macro- and microelements and optionally vitamins as active agents, which contain 20-50 % by mass of mannitol as shell material, 8-25 % by mass of potassium hydrogen carbonate as basic sparkling component, 9-27 % by mass of malic acid as acidic sparkling component, and 0.4-2.2 % by mass of aspartame as sweetening agent, furthermore possibly flavouring, lubricating and other additives generally used in the manufacture of effervescent tablets in an amount necessary to supplement the mass of the components to 100 %.

2. Effervescent tablets or granules as claimed in claim 1, which comprise 30-40 % by mass of mannitol, 14-18 % by mass of potassium hydrogen carbonate, 15-21 % by mass of malic acid, and 0.6-1.5 % by mass of aspartame.

3. Effervescent tablets or granules as claimed in claim 1, which contain as macro- and trace elements magnesium, zinc, iron(II), copper(II), manganese(II) and chromium(III) cations, further molybdenum(VI) and selenium(IV) anions.

4. Effervescent tablets or granules as claimed in claim 1, wherein the iron ions are contained in the form of ferro-sulphate heptahydrate, the zinc ions in the form of zinc sulphate heptahydrate, the copper ions in the form of copper sulphate pentahydrate, the manganese ions in the form of manganese sulphate monohydrate, the molybdenum ions in the form of ammonium heptamolybdenate tetrahydrate, the selenium ions in the form of selenious acid, the magnesium ions in the form of magnesium sulphate heptahydrate, and the chromium ions in the form of chromium(III) chloride hexahydrate.

5. Effervescent tablets or granules as claimed in claim 1, wherein the vitamins are present in the following amounts related to the mass of the composition: 0.01-0.5 % by mass of vitamin B<sub>1</sub>, 0.01-0.25 % by mass of vitamin B<sub>2</sub>, 0.01-0.5 % by mass of vitamin B<sub>6</sub>, 0.001-0.01 % by mass of vitamin B<sub>12</sub>,

- 14 -

0.1-2 % by mass of nicotinamide, 0.01-0.5 % by mass of vitamin A, 0.0015-0.015 % by mass of vitamin D, 0.1-5 % by mass of vitamin C, 0.01-0.1 % by mass of folic acid, 0.1-0.5% by mass of pantothenic acid, 0.01-7 % by mass of vitamin E and  
5 0.001-0.01 % by mass of vitamin H.

6. A process for preparing effervescent tablets or granules comprising the steps of homogenizing 20-50 % by mass of mannitol, 8-25 % by mass of potassium hydrogen carbonate, 9-24 % by mass of malic acid, and 0.4-2.2 % by mass of aspartame together with the macro- and trace elements and vitamins to be introduced and optionally together with flavouring, lubricating and other additives generally used in the manufacture of effervescent tablets, then granulating the thus-obtained homogenizate to granules ready for com-  
10 pressing, and finally compressing tablets or granules of the desired size and strength.

7. A process as claimed in claim 6, in which 30-40 % by mass of mannitol, 14-18 % by mass of potassium hydrogen carbonate, 15-21 % by mass of malic acid, and 0.6-1.5 % by  
20 mass of aspartame are homogenized with the other components of the tablet.



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/HU 94/00006

## A. CLASSIFICATION OF SUBJECT MATTER

IPC<sup>5</sup>: A 61 K 9/46; A 23 L 2/40; A 23 L 1/236; A 61 K 31/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC<sup>5</sup>: A 61 K 9/00; A 23 L 2/00; A 23 L 1/00; A 61 K 31/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP, A1, 0 396 335 (BEECHAM GROUP p.l.c.) 07 November 1990 (07.11.90), claims 1,2,4,7,8; abstract; examples 4C, 4D.	1,2,6,7
A	US, A, 4 127 645 (F. WITZEL et al.) 28 November 1978 (28.11.78), claims 1,3 to 8, 10-12; column 2, lines 10 to 52.	1,2,6,7
A	DE, A, 2 058 434 (ASPRO-NICHOLAS LTD.) 03 June 1971 (03.06.71), claims 1,2,15,17,18,22; example 1; page 10; line 8 to page 12, line 20.	1,2,6,7
A	US, A, 5 178 878 (F. WEHLING et al.) 12 January 1993 (12.01.93), abstract; column 4, line 56 to column 5, line 26; column 12, line 28 to column 13, line 15.	1-7

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;" document member of the same patent family

Date of the actual completion of the international search

06 July 1994 (06.07.94)

Date of mailing of the international search report

12 August 1994 (12.08.94)

Name and mailing address of the ISA/ AT  
AUSTRIAN PATENT OFFICE  
Kohlmarkt 8-10  
A-1014 Vienna  
Facsimile No. 1/53424/535

Authorized officer

Mazzucco e.h.

Telephone No. 1/5337058/33

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.

PCT/HU 94/00006

In Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
EP A1 396335	07-11-90	AT E 101034 AU A1 53914/90 AU B2 627298 CA AA 2015536 DE C0 69006385 DE T2 69006385 EP B1 396335 GB A0 8909793 JP A2 2295919 NZ A 233441 PT A 93877 US A 5225197 ZA A 9003181	15-02-94 01-11-90 20-08-92 28-10-90 17-03-94 19-05-94 02-02-94 14-06-89 06-12-90 26-03-92 20-11-90 06-07-93 26-06-91
US A 4127645	28-11-78	keine - none - rien	
DE A1 2058434		AT B 302533 BE A1 759520 CA A1 948108 DE A 2058434 DE B2 2058434 DE C3 2058434 DK B 130453 DK C 130453 ES A1 385974 FR A5 2073431 FR B1 2073431 GB A 1287475 IE B 34756 IL A0 35714 IL A1 35714 IN A 129401 JP B4 51006727 NL A 7017417 NL B 165928 NL C 165928 SE B 383099 SE C 383099 US A 3882228 US A 3887700 ZA A 7007915 ZM A 134/70	25-10-72 99-99-99 28-05-74 03-06-71 24-04-80 18-12-80 24-02-75 21-07-75 01-05-73 01-10-71 22-03-74 31-08-72 06-08-75 28-01-71 14-03-74 16-08-75 02-03-76 02-06-71 15-01-81 15-06-81 01-03-76 10-06-76 06-05-75 03-06-75 25-08-71 22-11-71
US A 5178878	12-01-93	AU A1 66012/90 AU B2 646232 EP A1 494972 EP A4 494972 JP T2 5500956 WO A1 9104757 US A 5223264 CA AA 2061917	28-04-91 17-02-94 22-07-92 12-08-92 25-02-93 18-04-91 29-06-93 27-08-93